

<b>POLICY TITLE</b>	<b>ACTIGRAPHY</b>
<b>POLICY NUMBER</b>	<b>MP 2.087</b>

<b>CLINICAL BENEFIT</b>	<input type="checkbox"/> MINIMIZE SAFETY RISK OR CONCERN. <input checked="" type="checkbox"/> MINIMIZE HARMFUL OR INEFFECTIVE INTERVENTIONS. <input type="checkbox"/> ASSURE APPROPRIATE LEVEL OF CARE. <input type="checkbox"/> ASSURE APPROPRIATE DURATION OF SERVICE FOR INTERVENTIONS. <input type="checkbox"/> ASSURE THAT RECOMMENDED MEDICAL PREREQUISITES HAVE BEEN MET. <input type="checkbox"/> ASSURE APPROPRIATE SITE OF TREATMENT OR SERVICE.
<b>Effective Date:</b>	<b>11/1/2024</b>

[POLICY RATIONALE](#)  
[DISCLAIMER](#)  
[POLICY HISTORY](#)

[PRODUCT VARIATIONS](#)  
[DEFINITIONS](#)  
[CODING INFORMATION](#)

[DESCRIPTION/BACKGROUND](#)  
[BENEFIT VARIATIONS](#)  
[REFERENCES](#)

**I. POLICY**

Actigraphy is considered **investigational** when used as the sole technique to record and analyze body movement, including but not limited to its use to evaluate sleep disorders. This does not include the use of actigraphy as a component of portable sleep monitoring. (See Policy Guideline section) There is insufficient evidence to support a general conclusion concerning the health outcomes or benefits associated with this procedure.

**Policy Guidelines**

This policy does not address the use of actigraphy as a component of portable sleep monitoring. When used as a component of portable sleep monitoring, actigraphy should be reported separately.

**\*Note:** The monitoring device necessary for this procedure will be provided by the physician conducting the examination.

**Cross-references:**

**MP 2.045** Diagnosis and Medical Management of Obstructive Sleep Apnea

**II. PRODUCT VARIATIONS**

[TOP](#)

This policy is only applicable to certain programs and products administered by Capital Blue Cross and subject to benefit variations as discussed in Section VI. Please see additional information below.

**FEP PPO-** Refer to FEP Medical Policy Manual. The FEP Medical Policy Manual can be found at: <https://www.fepblue.org/benefit-plans/medical-policies-and-utilization-management-guidelines/medical-policies>

<b>POLICY TITLE</b>	<b>ACTIGRAPHY</b>
<b>POLICY NUMBER</b>	<b>MP 2.087</b>

**III. DESCRIPTION/BACKGROUND**

[TOP](#)

Actigraphy refers to the assessment of body movement activity patterns using devices, typically placed on the wrist or ankle, during sleep, which are interpreted by computer algorithms as periods of sleep and wake. Sleep-wake cycles may be altered in sleep disorders, including insomnia and circadian rhythm sleep disorders. Also, actigraphy could be used to assess sleep/wake disturbances associated with other disorders.

**Sleep Disorders**

Sleep disorders affect a large percentage of the U.S. population. For example, estimates suggest that 15% to 24% of the U.S. population suffers from insomnia. Lack of sleep also contributes reduced cognitive functioning, susceptibility to heart disease, and workplace absenteeism.

**Diagnosis**

Actigraphy refers to the assessment of activity patterns (body movement) using devices, typically placed on the wrist or ankle, which are interpreted by computer algorithms as periods of sleep (absence of activity) and wake (activity). Actigraphy devices are usually placed on the nondominant wrist with a wristband and are worn continuously for at least 24 hours. Activity is usually recorded for a period of 3 days to 2 weeks but can be collected continuously over extended periods with regular downloading of data onto a computer. The activity monitors may also be placed on the ankle to assess restless legs syndrome or on the trunk to record movement in infants.

The algorithms for detecting movement vary across devices and may include “time above threshold,” the “zero crossing method” (the number of times per epoch that activity level crosses zero), or “digital integration” method, resulting in different sensitivities. Sensitivity settings (e.g., low, medium, high, automatic) can also be adjusted during data analysis. The most commonly used method (digital integration) reflects both acceleration and amplitude of movement.

Data on patient bedtimes (lights out) and rise times (lights on) are usually entered into the computer from daily patient sleep logs or by patient-activated event markers. Proprietary software is then used to calculate periods of sleep based on the absence of detectable movement, along with the movement related level of activity and periods of wake. In addition to providing a graphic depiction of the activity pattern, the device-specific software can then analyze and report a variety of sleep parameters, including sleep onset, sleep offset, sleep latency, total sleep duration, and wake after sleep onset (actigraphy could also be used to measure the level of physical activity).

Actigraphy has been used for more than two decades as an outcome measure in sleep disorders research. For clinical applications, actigraphy is being evaluated as a measure of sleep-wake cycles in sleep disorders, including insomnia and circadian rhythm sleep disorders. Also, actigraphy is being investigated as a measure of sleep-wake disturbances associated with other diseases and disorders.

**Regulatory Status**

Numerous actigraphy devices have been cleared for marketing by the U.S. Food and Drug Administration through the 510(k) process. Some actigraphy devices are designed and marketed

<b>POLICY TITLE</b>	<b>ACTIGRAPHY</b>
<b>POLICY NUMBER</b>	<b>MP 2.087</b>

to measure sleep-wake states while others to measure levels of physical activity. Food and Drug Administration product code: OLV.

**IV. RATIONALE**

[TOP](#)

**Summary of Evidence**

For individuals who have circadian sleep-wake rhythm disorders who receive actigraphy, the evidence includes a comparative study that selected subjects from another main study evaluating the effects of caffeine on daytime recovery sleep. Relevant outcomes are test accuracy and test validity. Comparison with polysomnography (PSG) has shown that actigraphy is limited in differentiating between sleep and wake in more disturbed sleep. Actigraphy appears to reliably measure sleep onset and total sleep time in some patient populations. Comparisons with PSG and sleep diaries are limited. Evidence has shown that actigraphy does not provide a reliable measure of sleep efficiency in this patient population. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For children and adolescents with sleep-associated disorders who receive actigraphy, the evidence includes prospective and retrospective validation studies. Relevant outcomes are test accuracy and validity. Comparisons with PSG have shown that actigraphy can differ significantly in its estimations of wake and sleep times and sleep onset latency. Comparisons with sleep diaries have also failed to show satisfactory agreement, with greater discrepancies for more disturbed sleep. Evidence has shown that actigraphy does not provide a reliable measure of sleep efficiency in this patient population. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have central disorders of hypersomnolence who receive actigraphy, the evidence includes a comparative observational study. Relevant outcomes are test accuracy and validity. Comparison with video-PSG has indicated that actigraphy has a sensitivity of 26.1% and specificity of 95.5%. General evidence has also revealed that the accuracy of actigraphy for differentiating between wake and sleep decreases as the level of sleep disturbance increases. Although actigraphy appears to provide reliable measures of sleep onset and wake time in some patient populations, its clinical utility compared with that of sleep diaries has not been demonstrated. Evidence has shown that actigraphy does not provide a reliable measure of sleep efficiency in this patient population. The complexity of the various syndromes as well as the potential for medical treatment with significant adverse events makes accurate diagnosis essential. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have insomnia who receive actigraphy, the evidence includes prospective and retrospective validation studies. Relevant outcomes are test accuracy and validity. Comparisons with PSG have shown that actigraphy has poor agreement for reporting wake time and can overestimate sleep efficiency. Comparison with sleep diaries has indicated that actigraphy is less effective at differentiating between patients with insomnia and controls. General evidence has also revealed that the accuracy of actigraphy for differentiating between wake and sleep decreases as the level of sleep disturbance increases. Although actigraphy appears to provide reliable measures of sleep onset and wake time in some patient populations, its clinical

<b>POLICY TITLE</b>	<b>ACTIGRAPHY</b>
<b>POLICY NUMBER</b>	<b>MP 2.087</b>

utility compared with sleep diaries has not been demonstrated. Evidence has shown that actigraphy does not provide a reliable measure of sleep efficiency in this patient population. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

**V. DEFINITIONS**

[TOP](#)

**510 (K)** is a premarketing submission made to FDA to demonstrate that the device to be marketed is as safe and effective, that is, substantially equivalent (SE), to a legally marketed device that is not subject to premarket approval (PMA). Applicants must compare their 510(k) device to one or more similar devices currently on the U.S. market and make and support their substantial equivalency claims.

**APNEA** is the cessation of respiration for at least ten (10) seconds.

**VI. BENEFIT VARIATIONS**

[TOP](#)

The existence of this medical policy does not mean that this service is a covered benefit under the member's health benefit plan. Benefit determinations should be based in all cases on the applicable health benefit plan language. Medical policies do not constitute a description of benefits. A member's health benefit plan governs which services are covered, which are excluded, which are subject to benefit limits, and which require preauthorization. There are different benefit plan designs in each product administered by Capital Blue Cross. Members and providers should consult the member's health benefit plan for information or contact Capital Blue Cross for benefit information.

**VII. DISCLAIMER**

[TOP](#)

*Capital Blue Cross's medical policies are developed to assist in administering a member's benefits, do not constitute medical advice and are subject to change. Treating providers are solely responsible for medical advice and treatment of members. Members should discuss any medical policy related to their coverage or condition with their provider and consult their benefit information to determine if the service is covered. If there is a discrepancy between this medical policy and a member's benefit information, the benefit information will govern. Capital Blue Cross considers the information contained in this medical policy to be proprietary and it may only be disseminated as permitted by law.*

**VIII. CODING INFORMATION**

[TOP](#)

**Note:** This list of codes may not be all-inclusive, and codes are subject to change at any time. The identification of a code in this section does not denote coverage as coverage is determined by the terms of member benefit information. In addition, not all covered services are eligible for separate reimbursement.

**Investigational; therefore, not covered:**

Procedure Codes								
95803								

<b>POLICY TITLE</b>	<b>ACTIGRAPHY</b>
<b>POLICY NUMBER</b>	<b>MP 2.087</b>

**IX. REFERENCES**

[TOP](#)

1. Ford ES, Cunningham TJ, Giles WH, et al. Trends in insomnia and excessive daytime sleepiness among U.S. adults from 2002 to 2012. *Sleep Med.* Mar 2015; 16(3): 372-8. PMID 25747141
2. Paquet J, Kawinska A, Carrier J. Wake detection capacity of actigraphy during sleep. *Sleep.* Oct 2007; 30(10): 1362-9. PMID 17969470
3. Meltzer LJ, Wong P, Biggs SN, et al. Validation of Actigraphy in Middle Childhood. *Sleep.* Jun 01, 2016; 39(6): 1219-24. PMID 27091520
4. Enomoto M, Kitamura S, Nakazaki K. Validity of an algorithm for determining sleep/wake states using FS-760 in school-aged children. *J Physiol Anthropol.* Aug 18 2022; 41(1): 29. PMID 35982481
5. Yavuz-Kodat E, Reynaud E, Geoffroy MM, et al. Validity of Actigraphy Compared to Polysomnography for Sleep Assessment in Children With Autism Spectrum Disorder. *Front Psychiatry.* 2019; 10: 551. PMID 31428003
6. O'Driscoll DM, Foster AM, Davey MJ, et al. Can actigraphy measure sleep fragmentation in children? *Arch Dis Child.* Dec 2010; 95(12): 1031-3. PMID 19850594
7. Hyde M, O'Driscoll DM, Binette S, et al. Validation of actigraphy for determining sleep and wake in children with sleep disordered breathing. *J Sleep Res.* Jun 2007; 16(2): 213-6. PMID 17542951
8. Bélanger MÈ, Bernier A, Paquet J, et al. Validating actigraphy as a measure of sleep for preschool children. *J Clin Sleep Med.* Jul 15, 2013; 9(7): 701-6. PMID 23853565
9. Insana SP, Gozal D, Montgomery-Downs HE. Invalidity of one actigraphy brand for identifying sleep and wake among infants. *Sleep Med.* Feb 2010; 11(2): 191-6. PMID 20083430
10. Spruyt K, Gozal D, Dayyat E, et al. Sleep assessments in healthy school-aged children using actigraphy: concordance with polysomnography. *J Sleep Res.* Mar 2011; 20(1 Pt 2): 223-32. PMID 20629939
11. Werner H, Molinari L, Guyer C, et al. Agreement rates between actigraphy, diary, and questionnaire for children's sleep patterns. *Arch Pediatr Adolesc Med.* Apr 2008; 162(4): 350-8. PMID 18391144
12. Short MA, Gradisar M, Lack LC, et al. The discrepancy between actigraphic and sleep diary measures of sleep-in adolescents. *Sleep Med.* Apr 2012; 13(4): 378-84. PMID 22437142
13. Louter M, Arends JB, Bloem BR, et al. Actigraphy as a diagnostic aid for REM sleep behavior disorder in Parkinson's disease. *BMC Neurol.* Apr 06, 2014; 14: 76. PMID 24708629
14. Marino M, Li Y, Rueschman MN, et al. Measuring sleep: accuracy, sensitivity, and specificity of wrist actigraphy compared to polysomnography. *Sleep.* Nov 01, 2013; 36(11): 1747-55. PMID 24179309
15. Taibi DM, Landis CA, Vitiello MV. Concordance of polysomnographic and actigraphic measurement of sleep and wake in older women with insomnia. *J Clin Sleep Med.* Mar 15, 2013; 9(3): 217-25. PMID 23493815
16. Levenson JC, Troxel WM, Begley A, et al. A quantitative approach to distinguishing older adults with insomnia from good sleeper controls. *J Clin Sleep Med.* Feb 01, 2013; 9(2): 125-31. PMID 23372464

<b>POLICY TITLE</b>	<b>ACTIGRAPHY</b>
<b>POLICY NUMBER</b>	<b>MP 2.087</b>

17. Kaplan KA, Talbot LS, Gruber J, et al. Evaluating sleep in bipolar disorder: comparison between actigraphy, polysomnography, and sleep diary. *Bipolar Disord.* Dec 2012; 14(8): 870-9. PMID 23167935
18. Dick R, Penzel T, Fietze I, et al. AASM standards of practice compliant validation of actigraphic sleep analysis from SOMNOWatch versus polysomnographic sleep diagnostics shows high conformity also among subjects with sleep disordered breathing. *Physiol Meas.* Dec 2010; 31(12): 1623-33. PMID 21071830
19. Sivertsen B, Omvik S, Havik OE, et al. A comparison of actigraphy and polysomnography in older adults treated for chronic primary insomnia. *Sleep.* Oct 2006; 29(10): 1353-8. PMID 17068990
20. Smith MT, McCrae CS, Cheung J, et al. Use of Actigraphy for the Evaluation of Sleep Disorders and Circadian Rhythm Sleep-Wake Disorders: An American Academy of Sleep Medicine Clinical Practice Guideline. *J Clin Sleep Med.* Jul 15, 2018; 14(7): 1231-1237. PMID 29991437
21. Blue Cross Blue Shield Association Medical Policy Reference Manual. 2.01.73, Actigraphy. July 2023

**X. POLICY HISTORY**

[TOP](#)

<b>MP 2.087</b>	<b>07/26/2011 CAC New policy</b> adopting BCBSA. Information related to Actigraphy was extracted from MP 2.045 Sleep Apnea Syndrome and a new policy was created. Policy statement unchanged.
	<b>01/29/2013 CAC Consensus review.</b> References updated. No changes to the policy statements. FEP variation revised to refer to the FEP medical policy manual. Codes reviewed 12/12/12
	<b>01/28/2014 CAC Consensus review.</b> No change to policy statements. References updated. Rationale section added.
	<b>01/27/2015 CAC Consensus review.</b> No change to the policy statement. References and rationale updated. No coding changes
	<b>01/26/2016 CAC Consensus review.</b> No change to the policy statement. References and rationale updated. Coding reviewed.
	<b>11/10/2016 Admin Update</b> -Variation reformatting
	<b>01/31/2017 – CAC Consensus review.</b> No change to policy statements. References and rationale reviewed. Coding reviewed
	<b>12/01/2017 Consensus review.</b> No change to policy statements. References and rationale updated.
	<b>08/28/2018 Consensus review.</b> Clarification to the policy statement this policy does not apply to actigraphy used for portable sleep monitoring. Background and references updated. Rationale revised.
	<b>06/14/2019 Consensus review.</b> No change to policy statements.

<b>POLICY TITLE</b>	<b>ACTIGRAPHY</b>
<b>POLICY NUMBER</b>	<b>MP 2.087</b>

<p><b>07/1/2020 Consensus review.</b> No change to the policy statement. References reviewed/updated. Coding reviewed with no changes. Product variation statement updated.</p>
<p><b>03/10/2021- Consensus review.</b> No change to the policy statement. References reviewed/updated. Coding reviewed with no changes.</p>
<p><b>09/02/2021 Administrative update.</b> Remove policy name and number from the product variation section.</p>
<p><b>07/12/2022 Consensus review.</b> No change to policy statement. References reviewed and updated. Coding reviewed with no changes.</p>
<p><b>07/11/2023 Consensus review.</b> No change to policy statement. Background and Rationale updated. References added.</p>
<p><b>1/18/2024 Administrative Update.</b> Added clinical benefit.</p>

[TOP](#)

*Health care benefit programs issued or administered by Capital Blue Cross and/or its subsidiaries, Capital Advantage Insurance Company®, Capital Advantage Assurance Company® and Keystone Health Plan® Central. Independent licensees of the Blue Cross BlueShield Association. Communications issued by Capital Blue Cross in its capacity as administrator of programs and provider relations for all companies.*