

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

CLINICAL BENEFIT	<input type="checkbox"/> MINIMIZE SAFETY RISK OR CONCERN. <input checked="" type="checkbox"/> MINIMIZE HARMFUL OR INEFFECTIVE INTERVENTIONS. <input type="checkbox"/> ASSURE APPROPRIATE LEVEL OF CARE. <input type="checkbox"/> ASSURE APPROPRIATE DURATION OF SERVICE FOR INTERVENTIONS. <input type="checkbox"/> ASSURE THAT RECOMMENDED MEDICAL PREREQUISITES HAVE BEEN MET. <input type="checkbox"/> ASSURE APPROPRIATE SITE OF TREATMENT OR SERVICE.
Effective Date:	6/1/2024

[POLICY RATIONALE](#)
[DISCLAIMER](#)
[POLICY HISTORY](#)

[PRODUCT VARIATIONS](#)
[DEFINITIONS](#)
[CODING INFORMATION](#)

[DESCRIPTION/BACKGROUND](#)
[BENEFIT VARIATIONS](#)
[REFERENCES](#)

I. POLICY

NOTE: Documentation of the usage of validated patient focused pain intensity assessment scales (such as Numeric Rating Scales, Visual Analog Scales, Verbal Rating Scales, the *Faces Pain Scale*, or the *Pain Thermometer*) is considered **medically necessary** in discriminating different levels of pain sensation during medical management.

Facet Joint Injections/Medial Branch Blocks

Initial Diagnostic Facet Joint Injection/Medial Branch Block

Facet joint injections/medial branch blocks may be considered **medically necessary** for lumbar or cervical pain lasting more than 3 months, despite appropriate conservative treatment, as a diagnostic procedure to determine whether the pain is of facet joint origin.

Second Diagnostic Facet Joint Injection/Medial Branch Block

A second diagnostic facet joint injection/medial branch block may be considered **medically necessary** when all the following criteria are met:

- The initial diagnostic injection was successful at relieving pain (see policy guidelines); AND
- The second diagnostic injection will be administered at the same level(s) as the initial diagnostic block; AND
- Radiofrequency joint denervation is being considered.

Therapeutic Facet Joint Injection/Medial Branch Block

Therapeutic facet joint injections/medial branch blocks may be considered **medically necessary** when all the following criteria are met:

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

- There is documented successful response with 2 sequential diagnostic facet joint injections/medial branch blocks at the same level(s); AND
- The individual is not a candidate for radiofrequency joint denervation; AND
- For subsequent therapeutic blocks there was successful response to the prior therapeutic injection for at least 3 months duration; AND
- For each spinal region, a maximum of 4 therapeutic facet joint sessions are medically necessary in a 12-month period.

Facet joint injection/medial branch block, diagnostic or therapeutic, for any indication other than the ones listed above is considered **investigational**. There is insufficient evidence to support a general conclusion concerning the health outcomes or benefits associated with this procedure.

Ultrasound guidance for facet joint injections is considered **investigational**. There is insufficient evidence to support a general conclusion concerning the health outcomes or benefits associated with this procedure.

Epidural Steroid Injections for Back Pain

Epidural steroid injections (transforaminal, caudal, interlaminar) performed with fluoroscopic guidance may be considered **medically necessary** for the treatment of pain when one of the following indications are met:

- Lumbar or cervical radiculopathy or sciatica that is not responsive to at least 4 weeks of conservative management (see Policy Guidelines section); **OR**
- Pain from herpes zoster or suspected radicular pain based on radiation of pain along the dermatome of a nerve; **OR**
- Pain from neurogenic claudication that includes any of the following:
 - Pain is severe enough to cause some degree of functional deficit
 - Failure of at least four weeks of noninvasive care
 - Imaging demonstrating a correlative region of nerve/cord impingement**OR**
- Back pain without lower extremities symptoms and failure of four weeks of non-surgical, non-injection care with either:
 - Documented Visual Analog Scale (VAS) for pain or Numeric Pain Rating Scale (NPRS) ≥3/10 (moderate to severe pain), **OR**
 - Functional impairment in ADLs; **AND**
 - The pain or functional impairment is associated with **any** of the following:
 - Substantial imaging abnormality, such as a central disc herniation or high intensity zone;
 - Documented severe degenerative disc disease or central spinal stenosis;
 - Discogenic pain, not attributable to facet joint or sacroiliac joint pain.

AND for any of the above medically necessary indications:

- Persistent pain is present of at least moderate-severe intensity; AND

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

- Short-term relief of pain is the anticipated outcome.

For transforaminal injections, simultaneous treatment of 2 contiguous vertebral levels (either unilateral or bilateral) performed during the same session may be considered **medically necessary** if criteria are met at each level.

- Note: when performing an interlaminar or caudal epidural injection, only 1 spinal level unilaterally is allowed during the same session.

For transforaminal injections, simultaneous treatment of more than 2 vertebral levels is considered **not medically necessary**.

Repeat treatment of persistent pain due to any of the above indications may be considered **medically necessary** under the following conditions:

- Previous epidural steroid injections were successful at relieving pain; AND
- At least 14 days have elapsed since the prior injection; AND
- For transforaminal, caudal or interlaminar steroid injections, no more than 6 injections per spinal region may be given over a 12-month period.

Repeat treatment is considered **not medically necessary** if the initial treatment did not result in substantial pain relief.

Epidural steroid injections are considered **investigational** in all other situations, including but not limited to nonspecific low back pain. There is insufficient evidence to support a general conclusion concerning the health outcomes or benefits associated with this procedure.

Epidural injection with ultrasound guidance for any indication is considered **investigational**, there is insufficient evidence to support a general conclusion concerning the health outcomes or benefits associated with this procedure.

The use of fluorography (imaging of the epidural space) as a component of epidural steroid injections is considered **investigational**. There is insufficient evidence to support a general conclusion concerning the health outcomes or benefits associated with this procedure.

POLICY GUIDELINES

Facet joint injection/medial branch block and epidural steroid injection by any approach should be performed with the use of fluoroscopic or CT guidance.

Facet Joint Injections/Medial Branch Blocks

Conservative therapies may include medications (e.g., non-steroidal anti-inflammatory drugs, antidepressants), physical treatments (exercise, heat or cold therapy, massage, physical therapy), integrative treatments (acupuncture, spinal manipulation if indicated) and others (nutrition, weight loss, sleep hygiene).

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

A successful block refers to a sustained meaningful reduction in pain. This is subjective but often defined as a 20-point improvement on a 0-to-100 visual analog scale or a 50% improvement using a validated scale.

Epidural Steroid Injections for Back Pain

The diagnosis of lumbar radiculopathy is typically made by a combination of suggestive signs and symptoms in conjunction with imaging that demonstrates compression of a spinal nerve root. Symptoms are due to irritation of the spinal nerve root at L4, L5, or S1, and may include posterior leg pain that extends past the knee, a loss of sensation in a dermatomal pattern, and/or loss of deep tendon reflexes. However, all of these symptoms may not be present. On exam, provocative tests such as the straight leg maneuver are positive. Magnetic resonance imaging is the most useful imaging modality and can confirm or exclude the presence of nerve root compression, most commonly due to herniated disc.

There are several aspects of epidural steroid injection therapy that are not standardized. Expert opinion was sought through clinical vetting on the following issues:

- The optimal time for assessing a response to epidural steroid injections. Expert opinion supports that response can be assessed anytime from immediately to several weeks after the procedure, with the most popular time to assess response being 1 to 2 weeks after injection.
- The definition of a clinically significant response to injections. Expert opinion supports that a reasonable definition of response is at least a 20-point improvement on a 0-to-100 visual analog scale, or an improvement of at least 50% in functional status, when measured using a validated scale.
- The maximum number of injections in 1 year. There is no agreement on the maximum number of injections that should be given in 1 year. Some experts agree that no more than 3 injections should be given in 1 year, but other experts believe that more than 3 per year can be used safely. None of the expert opinion supported more than 6 injections given over a 12-month period.

Conservative nonsurgical therapy for at least 4 weeks should include the following:

- Use of prescription-strength analgesics for several weeks at a dose sufficient to induce a therapeutic response.
 - Analgesics should include anti-inflammatory medications with or without adjunctive medications such as nerve membrane stabilizers or muscle relaxants AND
- Participation in at least 4 weeks of physical therapy (including active exercise) or documentation of why the individual could not tolerate physical therapy, AND
- Evaluation and appropriate management of associated cognitive and behavioral issues

Cross-reference:

- MP 2.072** Trigger Point and Tender Point Injections
- MP 4.041** Dry Needling of Myofascial Trigger Points

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

MP 5.048 Diagnosis and Treatment of Sacroiliac Joint Pain
MP 5.049 Facet Joint Denervation

II. PRODUCT VARIATIONS

[TOP](#)

This policy is only applicable to certain programs and products administered by Capital Blue Cross please see additional information below, and subject to benefit variations as discussed in Section VI below.

FEP PPO: Refer to FEP Medical Policy Manual. The FEP Medical Policy manual can be found at:

<https://www.fepblue.org/benefit-plans/medical-policies-and-utilization-management-guidelines/medical-policies>

III. DESCRIPTION/BACKGROUND

[TOP](#)

Back Pain

Back pain is an extremely common condition. Most episodes are self-limited and will resolve within 1 month, but a small percentage will persist and become chronic. Patients with chronic back pain may suffer from serious disability and may use a high volume of medical services. Despite high utilization, many patients with chronic back pain do not improve with available treatments, including surgical interventions. Therefore, there is a high unmet need to determine the efficacy of different treatments for chronic back pain and to determine which patient populations may benefit from specific interventions. In addition, there has been a proliferation of new technologies, combined with large increases in the number of patients treated and in the intensity of treatment. Therefore, there is a concern for overtreatment of patients who may not benefit from interventions for back pain.

Sciatica

Back pain can result from a variety of underlying causes. Sciatica is a subset of low back pain that is associated with irritation of one or more lumbar spinal nerve roots, which results in symptoms of radiculopathy. Symptoms of radiculopathy include pain that radiates down the leg to below the knee, numbness, muscle weakness, and lack of reflexes in a dermatomal distribution. Most patients with sciatica respond to conservative care with a resolution of their symptoms within several weeks to several months following onset. In a subset of patients, symptoms and signs of progressive muscle weakness prompt a more aggressive intervention to prevent permanent dysfunction. In other patients, symptoms persist despite conservative management, without progression of neurologic signs, and further treatment options are sought for pain relief.

Spinal Stenosis

Spinal stenosis is another common source of back pain. Spinal stenosis is caused by narrowing of the spinal canal due to degenerative changes, leading to impingement of the spinal cord and

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

the spinal nerve roots. Symptoms of spinal stenosis can include back pain, leg pain with exertion (neurogenic claudication), muscle weakness, and sensory deficits. Definitive treatment for spinal stenosis is surgery, which includes decompression of the spinal canal with or without spinal fusion. Epidural steroids may reduce inflammation from pressure on the spinal cord, and thus reduce symptoms of compression.

Nonspecific Low Back Pain

Nonspecific low back pain, sometimes called mechanical low back pain, is diagnosed when no specific etiology of pain can be identified. Although the etiology of nonspecific low back pain is uncertain, many experts feel that the pain is of discogenic origin or due to painful movement of the vertebrae. In these instances, epidural steroid injections may reduce swelling of the vertebral disc and/or surrounding structures, leading to pain relief.

Treatment

Regardless of specific etiology, conservative management is the first-line treatment for most patients with neck or back pain. Nonsteroidal anti-inflammatory drugs or other analgesics are used for symptom relief. These agents should be used for at least several weeks at a dose sufficient to induce a therapeutic response. Additionally, modification of activity in conjunction with some form of exercise therapy is frequently prescribed early in the course of symptoms and typically involves a physical therapist. For patients with persistent nonradicular back pain, current guidelines recommend interdisciplinary rehabilitation, which is defined as an integrated approach using physical rehabilitation in conjunction with a psychological or psychosocial intervention.

For patients who fail conservative therapy, a number of interventional therapies are available, which range from minimally invasive procedures, such as injections, to major surgeries, such as spinal decompression with fusion. Injections can be given in different locations (e.g., soft tissues, intraspinal, sacroiliac joints) and can use different therapeutic agents (e.g., botulinum toxin, steroids, proteolytic enzymes). Other interventional techniques include radiofrequency ablation, prolotherapy, and chemonucleolysis. Most of these nonsurgical interventions do not have high-quality evidence demonstrating their efficacy. A number of surgical interventions are available, such as discectomy and spinal fusion, each of which can be performed by a variety of techniques. The decision to undertake surgery is best made in the setting of shared decision making between the patient and surgeon, with thorough consideration given to the risks and benefits of surgery.

Epidural Steroid Injections

Epidural injection therapy is one of several second-line therapies available for patients who fail conservative treatment and is one of the most common modalities used in this group of patients. Epidural steroid injections are performed by inserting a needle into the space between the dura and ligamentum flavum and injecting a steroid preparation. There is considerable variability in the technical aspects of epidural injections. Several different approaches may be used for entering the epidural space (translaminar, transforaminal, caudal). In addition, epidural steroid injections may be administered with or without fluoroscopic guidance. For example, a national survey published in 2002 reported that 30% of academic institutions and 77% of private practices use fluoroscopy. Some investigators have estimated that lack of correct needle position

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

in the epidural space may occur in 25% or more of injections administered. Variability of the technique may also involve factors such as the depth of injection into the epidural space, the volume of injectate, and the filling patterns of the injectate.

Treatment is generally given as one to 3 injections, each performed at least 1 month apart. Some experts recommend no more than 3 injections in a 12-month period, owing to concerns about the adverse events of chronic steroid administration, both locally and systemically. Others contend that up to 6 injections per year are safe.

Facet Joint Injections/Medial Branch Blocks

Facet joint injections, also known as intra-articular injections (IA), are a minimally invasive procedure in which a physician injects a small amount of anesthetic, which may or may not contain a steroid, to numb the facet joint. An appropriately performed IA facet joint injection, in which the medication is only distributed to the joint capsule, can help to diagnosis facet joint pain. If the pain goes away after the IA injection, or is significantly decreased, then the facet joint can be determined to be the source of pain. While a good diagnostic tool for facet-mediated pain, IA is less predictive than medial branch blocks to medial branch radiofrequency ablation.

Medial branch blocks (MBB) are also used to diagnose facet joint pain and may or may not contain a steroid. MBB may not achieve the level of specificity that an appropriately performed IA injection can. This is due to the fact that the medial branch is not encapsulated and the medication can result in anesthesia of not only the medial branch, but also anesthesia of other nerves in close proximity. This could potentially result in non-selective neural blockade and a false-positive result. Even though this is a known possibility, MBB is an acceptable diagnostic test, and it is usually the preferred test when lumbar facet radiofrequency ablation is being considered. When used as a tool before lumbar facet neurolysis, MBB may be associated with a higher success rate than IA injections.

Once the facet joint has been determined to be the source of pain, a series of injections may be needed to establish consistency of results, especially if diagnostic facet joint injections/medial branch blocks are followed by neurolysis.

Image guidance has become an essential component of performing spinal procedures in pain management. For MBB and IA facet joint injections, fluoroscopy and to a lesser degree, CT guidance are most commonly used. The use of imaging allows accurate needle placement ensuring the lowest volume of anesthetic is administered, thereby reducing spread to surrounding tissues which may lead to false-positive results. Image guidance also improves safety as it helps to avoid unnecessary injury.

Regulatory Status

Steroids are not approved by the U.S. Food and Drug Administration for use as epidural injections; such use represents an off-label administration of a U.S. Food and Drug Administration-approved medication. The specific preparations used for epidural injections are steroids added to a sterile saline solution, which are prepared by a compounding pharmacy.

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

IV. RATIONALE

[Top](#)

Summary of Evidence

For individuals who have lumbar or cervical radiculopathy who receive ESI, the evidence includes many small RCTs and a number of systematic reviews of these RCTs. Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, medication use, and treatment-related morbidity. The evidence base lacks large-scale, high-quality trials and has a high degree of variability among the available trials in terms of patient populations, epidural injection techniques, and comparison treatments. The results of individual trials are mixed, with some reporting significant benefits for the ESI group and others reporting no benefit. Most systematic reviews did not perform pooled analyses due to the heterogeneity of trials. In the 2 reviews that reported quantitative results, short-term pain relief at up to 6 months follow-up was superior in patients treated with epidural steroids. None of the analyses reported long-term benefits for treatment with ESIs. Adverse events were generally mild but not well reported in these trials. Serious adverse events can occur, but their rate is unknown. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have spinal stenosis who receive ESIs, the evidence includes a moderately large RCT, a few small RCTs, and systematic reviews of these RCTs. Relevant outcomes include symptoms, functional outcomes, health status measures, quality of life, medication use, and treatment-related morbidity. The largest RCT and the majority of smaller trials did not report a benefit for ESIs. The evidence is insufficient to determine the effects of technology on health outcomes.

For individuals who have nonspecific low back pain who receive ESIs, the evidence includes a number of small RCTs and systematic reviews of these RCTs. Relevant outcomes include symptoms, functional outcomes, health status measures, quality of life, medication use, and treatment-related morbidity. Most trials were of low quality and did not report a benefit for ESIs. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have back or neck pain who receive facet joint injections/medial branch blocks, the clinical evidence for the efficacy of these injections are considered limited. Based on consensus practice guidelines and standards of care for those in chronic pain, the evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

V. DEFINITIONS

[Top](#)

ARTHROGRAPHY is a diagnostic study that involves the injection of contrast media into a joint.

FACET JOINT refers to one of the zygapophyseal joints of the vertebral column between the articulating facets of each pair of vertebrae.

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

INTRAVENOUS refers to within or into a vein.

NERVE BLOCK refers to interruption of the conduction of impulses to peripheral nerves or nerve trunks by the injection of a local anesthetic solution.

NUMERIC PAIN RATING SCALE- a pain assessment system in which patients are asked to rate their pain on a scale from 1 to 10, with 10 representing the worst pain they have experienced or could imagine.

OFF-LABEL USE: The use of a prescription drug or medical device in the treatment of an illness or injury for which it has not been specifically approved by the FDA.

PARAVERTEBRAL refers to alongside or near the vertebral column.

RADICULOPATHY refers to any disease of a nerve root.

VISUAL ANALOG SCALE (VAS) - A graphic scale that helps a patient to quantify pain, depression, and other subjective and otherwise unmeasurable states or conditions.

ZYGAPOPHYSEAL concerns a zygapophysis, one of the articular processes of a neural arch of a vertebrae.

VI. BENEFIT VARIATIONS

[TOP](#)

The existence of this medical policy does not mean that this service is a covered benefit under the member's health benefit plan. Benefit determinations should be based in all cases on the applicable health benefit plan language. Medical policies do not constitute a description of benefits. A member's health benefit plan governs which services are covered, which are excluded, which are subject to benefit limits, and which require preauthorization. There are different benefit plan designs in each product administered by Capital Blue Cross. Members and providers should consult the member's health benefit plan for information or contact Capital Blue Cross for benefit information.

VII. DISCLAIMER

[TOP](#)

Capital Blue Cross's medical policies are developed to assist in administering a member's benefits, do not constitute medical advice and are subject to change. Treating providers are solely responsible for medical advice and treatment of members. Members should discuss any medical policy related to their coverage or condition with their provider and consult their benefit information to determine if the service is covered. If there is a discrepancy between this medical policy and a member's benefit information, the benefit information will govern. If a provider or a member has a question concerning the application of this medical policy to a specific member's plan of benefits, please contact Capital Blue Cross' Provider Services or Member Services. Capital Blue Cross considers the information contained in this medical policy to be proprietary and it may only be disseminated as permitted by law.

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

VIII. CODING INFORMATION

[Top](#)

Note: This list of codes may not be all-inclusive, and codes are subject to change at any time. The identification of a code in this section does not denote coverage as coverage is determined by the terms of member benefit information. In addition, not all covered services are eligible for separate reimbursement.

Investigational; therefore, not covered, ultrasound guidance for facet joint injections:

Procedure Codes							
0213T	0214T	0215T	0216T	0217T	0218T		

Covered when medically necessary, facet joint nerve block injections:

Procedure Codes							
64490	64491	64492	64493	64494	64495		

ICD-10-CM Diagnosis Codes	Description
G54.1	Lumbosacral plexus disorders
G54.2	Cervical root disorders, not elsewhere classified
G54.3	Thoracic root disorders, not elsewhere classified
G54.4	Lumbosacral root disorders, not elsewhere classified
M08.1	Juvenile ankylosing spondylitis
M43.22	Fusion of spine, cervical region
M43.23	Fusion of spine, cervicothoracic region
M43.24	Fusion of spine, thoracic region
M43.25	Fusion of spine, thoracolumbar region
M43.26	Fusion of spine, lumbar region
M43.27	Fusion of spine, lumbosacral region
M43.28	Fusion of spine, sacral and sacrococcygeal region
M45.0	Ankylosing spondylitis of multiple sites in spine
M45.2	Ankylosing spondylitis of cervical region
M45.3	Ankylosing spondylitis of cervicothoracic region
M45.4	Ankylosing spondylitis of thoracic region
M45.5	Ankylosing spondylitis of thoracolumbar region
M45.6	Ankylosing spondylitis lumbar region
M45.7	Ankylosing spondylitis of lumbosacral region

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

ICD-10-CM Diagnosis Codes	Description
M45.8	Ankylosing spondylitis sacral and sacrococcygeal region
M45.A0	Non-radiographic axial spondyloarthritis of unspecified sites in spine
M45.A1	Non-radiographic axial spondyloarthritis of occipito-atlanto-axial region
M45.A2	Non-radiographic axial spondyloarthritis of cervical region
M45.A3	Non-radiographic axial spondyloarthritis of cervicothoracic region
M45.A4	Non-radiographic axial spondyloarthritis of thoracic region
M45.A5	Non-radiographic axial spondyloarthritis of thoracolumbar region
M45.A6	Non-radiographic axial spondyloarthritis of lumbar region
M45.A7	Non-radiographic axial spondyloarthritis of lumbosacral region
M45.A8	Non-radiographic axial spondyloarthritis of sacral and sacrococcygeal region
M45.AB	Non-radiographic axial spondyloarthritis of multiple sites in spine
M46.02	Spinal enthesopathy, cervical region
M46.03	Spinal enthesopathy, cervicothoracic region
M46.04	Spinal enthesopathy, thoracic region
M46.05	Spinal enthesopathy, thoracolumbar region
M46.06	Spinal enthesopathy, lumbar region
M46.07	Spinal enthesopathy, lumbosacral region
M46.08	Spinal enthesopathy, sacral and sacrococcygeal region
M46.09	Spinal enthesopathy, multiple sites in spine
M46.1	Sacroilitis, not elsewhere classified
M47.012	Anterior spinal artery compression syndromes, cervical region
M47.013	Anterior spinal artery compression syndromes, cervicothoracic region
M47.014	Anterior spinal artery compression syndromes, thoracic region
M47.015	Anterior spinal artery compression syndromes, thoracolumbar region
M47.016	Anterior spinal artery compression syndromes, lumbar region
M47.022	Vertebral artery compression syndromes, cervical region
M47.12	Other spondylosis with myelopathy, cervical region
M47.13	Other spondylosis with myelopathy, cervicothoracic region
M47.14	Other spondylosis with myelopathy, thoracic region
M47.15	Other spondylosis with myelopathy, thoracolumbar region
M47.16	Other spondylosis with myelopathy, lumbar region
M47.22	Other spondylosis with radiculopathy, cervical region
M47.23	Other spondylosis with radiculopathy, cervicothoracic region
M47.24	Other spondylosis with radiculopathy, thoracic region
M47.25	Other spondylosis with radiculopathy, thoracolumbar region

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

ICD-10-CM Diagnosis Codes	Description
M47.26	Other spondylosis with radiculopathy, lumbar region
M47.27	Other spondylosis with radiculopathy, lumbosacral region
M47.28	Other spondylosis with radiculopathy, sacral and sacrococcygeal region
M47.812	Spondylosis without myelopathy or radiculopathy, cervical region
M47.813	Spondylosis without myelopathy or radiculopathy, cervicothoracic region
M47.814	Spondylosis without myelopathy or radiculopathy, thoracic region
M47.815	Spondylosis without myelopathy or radiculopathy, thoracolumbar region
M47.816	Spondylosis without myelopathy or radiculopathy, lumbar region
M47.817	Spondylosis without myelopathy or radiculopathy, lumbosacral region
M47.818	Spondylosis without myelopathy or radiculopathy, sacral and sacrococcygeal region
M47.892	Other spondylosis, cervical region
M47.893	Other spondylosis, cervicothoracic region
M47.894	Other spondylosis, thoracic region
M47.895	Other spondylosis, thoracolumbar region
M47.896	Other spondylosis, lumbar region
M47.897	Other spondylosis, lumbosacral region
M47.898	Other spondylosis, sacral and sacrococcygeal region
M48.02	Spinal stenosis, cervical region
M48.03	Spinal stenosis, cervicothoracic region
M48.04	Spinal stenosis, thoracic region
M48.05	Spinal stenosis, thoracolumbar region
M48.061	Spinal stenosis, lumbar region without neurogenic claudication
M48.062	Spinal stenosis, lumbar region with neurogenic claudication
M48.07	Spinal stenosis, lumbosacral region
M48.08	Spinal stenosis, sacral and sacrococcygeal region
M48.8X2	Other specified spondylopathies, cervical region
M48.8X3	Other specified spondylopathies, cervicothoracic region
M48.8X4	Other specified spondylopathies, thoracic region
M48.8X5	Other specified spondylopathies, thoracolumbar region
M48.8X6	Other specified spondylopathies, lumbar region
M48.8X7	Other specified spondylopathies, lumbosacral region
M48.8X8	Other specified spondylopathies, sacral and sacrococcygeal region
M50.01	Cervical disc disorder with myelopathy, high cervical region
M50.021	Cervical disc disorder at C4-C5 level with myelopathy

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

ICD-10-CM Diagnosis Codes	Description
M50.022	Cervical disc disorder at C5-C6 level with myelopathy
M50.023	Cervical disc disorder at C6-C7 level with myelopathy
M50.03	Cervical disc disorder with myelopathy, cervicothoracic region
M50.11	Cervical disc disorder with radiculopathy, high cervical region
M50.121	Cervical disc disorder at C4-C5 level with radiculopathy
M50.122	Cervical disc disorder at C5-C6 level with radiculopathy
M50.123	Cervical disc disorder at C6-C7 level with radiculopathy
M50.13	Cervical disc disorder with radiculopathy, cervicothoracic region
M50.21	Other cervical disc displacement, high cervical region
M50.221	Other cervical disc displacement at C4-C5 level
M50.222	Other cervical disc displacement at C5-C6 level
M50.223	Other cervical disc displacement at C6-C7 level
M50.23	Other cervical disc displacement, cervicothoracic region
M50.31	Other cervical disc degeneration, high cervical region
M50.321	Other cervical disc degeneration at C4-C5 level
M50.322	Other cervical disc degeneration at C5-C6 level
M50.323	Other cervical disc degeneration at C6-C7 level
M50.33	Other cervical disc degeneration, cervicothoracic region
M50.81	Other cervical disc disorders, high cervical region
M50.821	Other cervical disc disorders at C4-C5 level
M50.822	Other cervical disc disorders at C5-C6 level
M50.823	Other cervical disc disorders at C6-C7 level
M50.83	Other cervical disc disorders, cervicothoracic region
M51.04	Intervertebral disc disorders with myelopathy, thoracic region
M51.05	Intervertebral disc disorders with myelopathy, thoracolumbar region
M51.06	Intervertebral disc disorders with myelopathy, lumbar region
M51.14	Intervertebral disc disorders with radiculopathy, thoracic region
M51.15	Intervertebral disc disorders with radiculopathy, thoracolumbar region
M51.16	Intervertebral disc disorders with radiculopathy, lumbar region
M51.17	Intervertebral disc disorders with radiculopathy, lumbosacral region
M51.24	Other intervertebral disc displacement, thoracic region
M51.25	Other intervertebral disc displacement, thoracolumbar region
M51.26	Other intervertebral disc displacement, lumbar region
M51.27	Other intervertebral disc displacement, lumbosacral region
M51.34	Other intervertebral disc degeneration, thoracic region

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

ICD-10-CM Diagnosis Codes	Description
M51.35	Other intervertebral disc degeneration, thoracolumbar region
M51.36	Other intervertebral disc degeneration, lumbar region
M51.37	Other intervertebral disc degeneration, lumbosacral region
M51.84	Other intervertebral disc disorders, thoracic region
M51.85	Other intervertebral disc disorders, thoracolumbar region
M51.86	Other intervertebral disc disorders, lumbar region
M51.87	Other intervertebral disc disorders, lumbosacral region
M53.2X7	Spinal instabilities, lumbosacral region
M53.2X8	Spinal instabilities, sacral and sacrococcygeal region
M53.3	Sacrococcygeal disorders, not elsewhere classified
M53.82	Other specified dorsopathies, cervical region
M53.83	Other specified dorsopathies, cervicothoracic region
M53.84	Other specified dorsopathies, thoracic region
M53.85	Other specified dorsopathies, thoracolumbar region
M53.86	Other specified dorsopathies, lumbar region
M53.87	Other specified dorsopathies, lumbosacral region
M53.88	Other specified dorsopathies, sacral and sacrococcygeal region
M54.2	Cervicalgia
M54.31	Sciatica, right side
M54.32	Sciatica, left side
M54.41	Lumbago with sciatica, right side
M54.42	Lumbago with sciatica, left side
M54.5	Low back pain
M54.50	Low back pain, unspecified
M54.51	Vertebrogenic low back pain
M54.59	Other low back pain
M54.6	Pain in thoracic spine
M54.89	Other dorsalgia
M62.830	Muscle spasm of back
M96.1	Postlaminectomy syndrome, not elsewhere classified
M99.21	Subluxation stenosis of neural canal of cervical region
M99.22	Subluxation stenosis of neural canal of thoracic region
M99.23	Subluxation stenosis of neural canal of lumbar region
M99.24	Subluxation stenosis of neural canal of sacral region
M99.31	Osseous stenosis of neural canal of cervical region

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

ICD-10-CM Diagnosis Codes	Description
M99.32	Osseous stenosis of neural canal of thoracic region
M99.33	Osseous stenosis of neural canal of lumbar region
M99.34	Osseous stenosis of neural canal of sacral region
M99.41	Connective tissue stenosis of neural canal of cervical region
M99.42	Connective tissue stenosis of neural canal of thoracic region
M99.43	Connective tissue stenosis of neural canal of lumbar region
M99.44	Connective tissue stenosis of neural canal of sacral region
M99.51	Intervertebral disc stenosis of neural canal of cervical region
M99.52	Intervertebral disc stenosis of neural canal of thoracic region
M99.53	Intervertebral disc stenosis of neural canal of lumbar region
M99.54	Intervertebral disc stenosis of neural canal of sacral region
M99.61	Osseous and spondylosis stenosis of intervertebral foramina of cervical region
M99.62	Osseous and spondylosis stenosis of intervertebral foramina of thoracic region
M99.63	Osseous and spondylosis stenosis of intervertebral foramina of lumbar region
M99.64	Osseous and spondylosis stenosis of intervertebral foramina of sacral region
M99.71	Connective tissue and disc stenosis of intervertebral foramina of cervical region
M99.72	Connective tissue and disc stenosis of intervertebral foramina of thoracic region
M99.73	Connective tissue and disc stenosis of intervertebral foramina of lumbar region
M99.74	Connective tissue and disc stenosis of intervertebral foramina of sacral region
S13.4XXA	Sprain of ligaments of cervical spine, initial encounter
S13.4XXD	Sprain of ligaments of cervical spine, subsequent encounter
S13.8XXA	Sprain of joints and ligaments of other parts of neck, initial encounter
S13.8XXD	Sprain of joints and ligaments of other parts of neck, subsequent encounter
S14.2XXA	Injury of nerve root of cervical spine, initial encounter
S14.2XXD	Injury of nerve root of cervical spine, subsequent encounter
S16.1XXA	Strain of muscle, fascia and tendon at neck level, initial encounter
S16.1XXD	Strain of muscle, fascia and tendon at neck level, subsequent encounter
S23.3XXA	Sprain of ligaments of thoracic spine, initial encounter
S23.3XXD	Sprain of ligaments of thoracic spine, subsequent encounter
S24.2XXA	Injury of nerve root of thoracic spine, initial encounter
S24.2XXD	Injury of nerve root of thoracic spine, subsequent encounter

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

ICD-10-CM Diagnosis Codes	Description
S33.5XXA	Sprain of ligaments of lumbar spine, initial encounter
S33.5XXD	Sprain of ligaments of lumbar spine, subsequent encounter
S33.6XXA	Sprain of sacroiliac joint, initial encounter
S33.6XXD	Sprain of sacroiliac joint, subsequent encounter
S33.8XXA	Sprain of other parts of lumbar spine and pelvis, initial encounter
S33.8XXD	Sprain of other parts of lumbar spine and pelvis, subsequent encounter
S34.21XA	Injury of nerve root of lumbar spine, initial encounter
S34.21XD	Injury of nerve root of lumbar spine, subsequent encounter
S34.22XA	Injury of nerve root of sacral spine, initial encounter
S34.22XD	Injury of nerve root of sacral spine, subsequent encounter
S34.4XXA	Injury of lumbosacral plexus, initial encounter
S34.4XXD	Injury of lumbosacral plexus, subsequent encounter

Covered when medically necessary, epidural steroid injections:

Procedure Codes							
62320	62321	62322	62323	64479	64480	64483	64484

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

ICD-10-CM Diagnosis Codes	Description
B02.23	Postherpetic polyneuropathy
B02.7	Disseminated zoster
B02.8	Zoster with other complications
B02.9	Zoster without complications
G54.4	Lumbosacral root disorders, not elsewhere classified
G89.18	Other acute postprocedural pain
G96.12	Meningeal adhesions (cerebral) (spinal)
G96.19	Other disorders of meninges, not elsewhere classified
G97.1	Other reaction to spinal and lumbar puncture
M43.12	Spondylolisthesis, cervical region
M43.13	Spondylolisthesis, cervicothoracic region
M43.14	Spondylolisthesis, thoracic region
M43.15	Spondylolisthesis, thoracolumbar region
M43.16	Spondylolisthesis, lumbar region
M43.17	Spondylolisthesis, lumbosacral region
M47.22	Other spondylosis with radiculopathy, cervical region
M47.23	Other spondylosis with radiculopathy, cervicothoracic region
M47.24	Other spondylosis with radiculopathy, thoracic region
M47.25	Other spondylosis with radiculopathy, thoracolumbar region
M47.26	Other spondylosis with radiculopathy, lumbar region
M47.27	Other spondylosis with radiculopathy, lumbosacral region
M47.812	Spondylosis without myelopathy or radiculopathy, cervical region
M47.813	Spondylosis without myelopathy or radiculopathy, cervicothoracic region
M47.814	Spondylosis without myelopathy or radiculopathy, thoracic region
M47.815	Spondylosis without myelopathy or radiculopathy, thoracolumbar region
M47.816	Spondylosis without myelopathy or radiculopathy, lumbar region
M47.817	Spondylosis without myelopathy or radiculopathy, lumbosacral region
M48.02	Spinal stenosis, cervical region
M48.03	Spinal stenosis, cervicothoracic region
M48.04	Spinal stenosis, thoracic region
M48.05	Spinal stenosis, thoracolumbar region
M48.062	Spinal stenosis, lumbar region with neurogenic claudication
M48.07	Spinal stenosis, lumbosacral region
M50.121	Cervical disc disorder at C4-C5 level with radiculopathy
M50.122	Cervical disc disorder at C5-C6 level with radiculopathy

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

M50.123	Cervical disc disorder at C6-C7 level with radiculopathy
M50.13	Cervical disc disorder with radiculopathy, cervicothoracic region
M50.221	Other cervical disc displacement at C4-C5 level
M50.222	Other cervical disc displacement at C5-C6 level
M50.223	Other cervical disc displacement at C6-C7 level
M50.31	Other cervical disc degeneration, high cervical region
M51.14	Intervertebral disc disorders with radiculopathy, thoracic region
M51.15	Intervertebral disc disorders with radiculopathy, thoracolumbar region
M51.16	Intervertebral disc disorders with radiculopathy, lumbar region
M51.17	Intervertebral disc disorders with radiculopathy, lumbosacral region
M51.24	Other intervertebral disc displacement, thoracic region
M51.25	Other intervertebral disc displacement, thoracolumbar region
M51.26	Other intervertebral disc displacement, lumbar region
M51.27	Other intervertebral disc displacement, lumbosacral region
M51.34	Other intervertebral disc degeneration, thoracic region
M51.35	Other intervertebral disc degeneration, thoracolumbar region
M51.36	Other intervertebral disc degeneration, lumbar region
M51.37	Other intervertebral disc degeneration, lumbosacral region
M53.3	Sacrococcygeal disorders, not elsewhere classified
M53.86	Other specified dorsopathies, lumbar region
M53.87	Other specified dorsopathies, lumbosacral region
M54.2	Cervicalgia
M54.12	Radiculopathy, cervical region
M54.13	Radiculopathy, cervicothoracic region
M54.14	Radiculopathy, thoracic region
M54.15	Radiculopathy, thoracolumbar region
M54.16	Radiculopathy, lumbar region
M54.17	Radiculopathy, lumbosacral region
M54.18	Radiculopathy, sacral and sacrococcygeal region
M54.31	Sciatica, right side
M54.32	Sciatica, left side
M54.41	Lumbago with sciatica, right side
M54.42	Lumbago with sciatica, left side
M54.5	Low back pain
M54.50	Low back pain, unspecified
M54.51	Vertebrogenic low back pain
M54.59	Other low back pain

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

M54.6	Pain in thoracic spine
M96.1	Postlaminectomy syndrome, not elsewhere classified
M99.21	Subluxation stenosis of neural canal of cervical region
M99.22	Subluxation stenosis of neural canal of thoracic region
M99.23	Subluxation stenosis of neural canal of lumbar region
M99.24	Subluxation stenosis of neural canal of sacral region
M99.31	Osseous stenosis of neural canal of cervical region
M99.32	Osseous stenosis of neural canal of thoracic region
M99.33	Osseous stenosis of neural canal of lumbar region
M99.34	Osseous stenosis of neural canal of sacral region
M99.41	Connective tissue stenosis of neural canal of cervical region
M99.42	Connective tissue stenosis of neural canal of thoracic region
M99.43	Connective tissue stenosis of neural canal of lumbar region
M99.44	Connective tissue stenosis of neural canal of sacral region
M99.51	Intervertebral disc stenosis of neural canal of cervical region
M99.52	Intervertebral disc stenosis of neural canal of thoracic region
M99.53	Intervertebral disc stenosis of neural canal of lumbar region
M99.61	Osseous and subluxation stenosis of intervertebral foramina of cervical region
M99.62	Osseous and subluxation stenosis of intervertebral foramina of thoracic region
M99.63	Osseous and subluxation stenosis of intervertebral foramina of lumbar region
M99.71	Connective tissue and disc stenosis of intervertebral foramina of cervical region
M99.72	Connective tissue and disc stenosis of intervertebral foramina of thoracic region
M99.73	Connective tissue and disc stenosis of intervertebral foramina of lumbar region

IX. REFERENCES

[TOP](#)

Epidural Steroid Injections for Back Pain

1. Chou R, Loeser JD, Owens DK, et al. *Interventional therapies, surgery, and interdisciplinary rehabilitation for low back pain: an evidence-based clinical practice guideline from the American Pain Society. Spine (Phila Pa 1976). May 01 2009; 34(10): 1066-77. PMID 19363457*

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

2. *Benyamin RM, Manchikanti L, Parr AT, et al. The effectiveness of lumbar interlaminar epidural injections in managing chronic low back and lower extremity pain. Pain Physician. Jul-Aug 2012; 15(4): E363-404. PMID 22828691*
3. *Pinto RZ, Maher CG, Ferreira ML, et al. Epidural corticosteroid injections in the management of sciatica: a systematic review and meta-analysis. Ann Intern Med. Dec 18 2012; 157(12): 865-77. PMID 23362516*
4. *Qaseem A, Wilt TJ, McLean RM, et al. Noninvasive Treatments for Acute, Subacute, and Chronic Low Back Pain: A Clinical Practice Guideline From the American College of Physicians. Ann Intern Med. Apr 04 2017; 166(7): 514-530. PMID 28192789*
5. *Chou R, Atlas SJ, Stanos SP, et al. Nonsurgical interventional therapies for low back pain: a review of the evidence for an American Pain Society clinical practice guideline. Spine (Phila Pa 1976). May 01 2009; 34(10): 1078-93. PMID 19363456*
6. *Manchikanti L, Buenaventura RM, Manchikanti KN, et al. Effectiveness of therapeutic lumbar transforaminal epidural steroid injections in managing lumbar spinal pain. Pain Physician. May-Jun 2012; 15(3): E199-245. PMID 22622912*
7. *Verheijen EJA, Bonke CA, Amorij EMJ, et al. Epidural steroid compared to placebo injection in sciatica: a systematic review and meta-analysis. Eur Spine J. Nov 2021; 30(11): 3255-3264. PMID 33974132*
8. *Yang S, Kim W, Kong HH, et al. Epidural steroid injection versus conservative treatment for patients with lumbosacral radicular pain: A meta-analysis of randomized controlled trials. Medicine (Baltimore). Jul 24 2020; 99(30): e21283. PMID 32791709*
9. *Smith CC, McCormick ZL, Mattie R, et al. The Effectiveness of Lumbar Transforaminal Injection of Steroid for the Treatment of Radicular Pain: A Comprehensive Review of the Published Data. Pain Med. Mar 01 2020; 21(3): 472-487. PMID 31343693*
10. *Arirachakaran A, Siripaiboonkij M, Pairuchvej S, et al. Comparative outcomes of epidural steroids versus placebo after lumbar discectomy in lumbar disc herniation: a systematic review and meta-analysis of randomized controlled trials. Eur J Orthop Surg Traumatol. Dec 2018; 28(8): 1589-1599. PMID 29845327*
11. *Bhatia A, Flamer D, Shah PS, et al. Transforaminal Epidural Steroid Injections for Treating Lumbosacral Radicular Pain from Herniated Intervertebral Discs: A Systematic Review and Meta-Analysis. Anesth Analg. Mar 2016; 122(3): 857-870. PMID 26891397*
12. *Borton ZM, Oakley BJ, Clamp JA, et al. Cervical transforaminal epidural steroid injections for radicular pain : a systematic review. Bone Joint J. May 2022; 104-B(5): 567-574. PMID 35491579*
13. *Diwan S, Manchikanti L, Benyamin RM, et al. Effectiveness of cervical epidural injections in the management of chronic neck and upper extremity pain. Pain Physician. Jul-Aug 2012; 15(4): E405-34. PMID 22828692*
14. *Benyamin RM, Singh V, Parr AT, et al. Systematic review of the effectiveness of cervical epidurals in the management of chronic neck pain. Pain Physician. Jan-Feb 2009; 12(1): 137-57. PMID 19165300*
15. *Cohen SP, Hayek S, Semenov Y, et al. Epidural steroid injections, conservative treatment, or combination treatment for cervical radicular pain: a multicenter, randomized, comparative-effectiveness study. Anesthesiology. Nov 2014; 121(5): 1045-55. PMID 25335172*

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

16. Schneider MJ, Ammendolia C, Murphy DR, et al. Comparative Clinical Effectiveness of Nonsurgical Treatment Methods in Patients With Lumbar Spinal Stenosis: A Randomized Clinical Trial. *JAMA Netw Open*. Jan 04 2019; 2(1): e186828. PMID 30646197
17. Friedly JL, Comstock BA, Turner JA, et al. A randomized trial of epidural glucocorticoid injections for spinal stenosis. *N Engl J Med*. Jul 03 2014; 371(1): 11-21. PMID 24988555
18. Friedly JL, Comstock BA, Turner JA, et al. Long-Term Effects of Repeated Injections of Local Anesthetic With or Without Corticosteroid for Lumbar Spinal Stenosis: A Randomized Trial. *Arch Phys Med Rehabil*. Aug 2017; 98(8): 1499-1507.e2. PMID 28396242
19. Staal JB, de Bie R, de Vet HC, et al. Injection therapy for subacute and chronic low-back pain. *Cochrane Database Syst Rev*. Jul 16 2008; (3): CD001824. PMID 18646078
20. Koes BW, Scholten RJPM, Mens JMA, et al. Efficacy of epidural steroid injections for low-back pain and sciatica: a systematic review of randomized clinical trials. *Pain*. Dec 1995; 63(3): 279-288. PMID 8719528
21. Rathmell JP, Benzon HT, Dreyfuss P, et al. Safeguards to prevent neurologic complications after epidural steroid injections: consensus opinions from a multidisciplinary working group and national organizations. *Anesthesiology*. May 2015; 122(5): 974-84. PMID 25668411
22. US Food and Drug Administration. Epidural Corticosteroid Injection: Drug Safety Communication - Risk of Rare But Serious Neurologic Problems. 2014
23. Centers for Disease Control and Prevention (CDC). Multistate outbreak of fungal infection associated with injection of methylprednisolone acetate solution from a single compounding pharmacy - United States, 2012. *MMWR Morb Mortal Wkly Rep*. Oct 19 2012; 61(41): 839-42. PMID 23076093
24. Bise S, Pesquer L, Feldis M, et al. Comparison of three CT-guided epidural steroid injection approaches in 104 patients with cervical radicular pain: transforaminal anterolateral, posterolateral, and transfacet indirect. *Skeletal Radiol*. Dec 2018; 47(12): 1625-1633. PMID 30032466
25. American Academy of Neurology (AAN). AAN Summary of Evidence-Based Guidelines for Clinicians: Use of epidural steroid injections to treat lumbosacral radicular pain. 2007 (Reaffirmed on April 30, 2022).
26. Watters WC, Resnick DK, Eck JC, et al. Guideline update for the performance of fusion procedures for degenerative disease of the lumbar spine. Part 13: injection therapies, low-back pain, and lumbar fusion. *J Neurosurg Spine*. Jul 2014; 21(1): 79-90. PMID 24980590
27. Chou R, Qaseem A, Snow V, et al. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society. *Ann Intern Med*. Oct 02 2007; 147(7): 478-91. PMID 17909209
28. Benzon HT, Connis RT, De Leon-Casasola OA, et al. Practice guidelines for chronic pain management: an updated report by the American Society of Anesthesiologists Task Force on Chronic Pain Management and the American Society of Regional Anesthesia and Pain Medicine. *Anesthesiology*. Apr 2010; 112(4): 810-33. PMID 20124882
29. Manchikanti L, Knezevic NN, Navani A, et al. Epidural Interventions in the Management of Chronic Spinal Pain: American Society of Interventional Pain Physicians (ASIPP)

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

Comprehensive Evidence-Based Guidelines. Pain Physician. Jan 2021; 24(S1): S27-S208. PMID 33492918

30. *Sayed D, Grider J, Strand N, et al. The American Society of Pain and Neuroscience (ASPN) Evidence-Based Clinical Guideline of Interventional Treatments for Low Back Pain. J Pain Res. 2022; 15: 3729-3832. PMID 36510616*
31. *Deer TR, Grider JS, Pope JE, et al. Best Practices for Minimally Invasive Lumbar Spinal Stenosis Treatment 2.0 (MIST): Consensus Guidance from the American Society of Pain and Neuroscience (ASPN). J Pain Res. 2022; 15: 1325-1354. PMID 35546905*
32. *North American Spine Society (NASS). Evidence-based clinical guidelines for multidisciplinary spine care: Diagnosis and treatment of low back pain. 2020*
33. *Blue Cross Blue Shield Association Medical Policy Reference Manual. 2.01.94, Epidural Steroid Injections for Back Pain. December 2023.*

Facet Nerve Blocks

1. *American College of Occupational and Environmental Medicine. Occupational Medicine Practice Guidelines: Evaluation and Management of Common Health Problems and Functional Recovery of Workers. 2nd Ed. Beverly Farms, MA: OEM Press; 2004: pp. 300 and 309.*
2. *Boswell MV, Colson JD, Sehgal N, Dunbar EE, et al. A systematic review of therapeutic facet joint interventions in chronic spinal pain. Pain Physician 2007; 10: 229-253.*
3. *Boswell MV, Trescot AM, Datta S, et al. Interventional techniques: evidence-based practice guidelines in the management of chronic spinal pain. Pain Physician 2007; 10: 7-111.*
4. *Capital BlueCross Integrated Low Back Pain Clinical Practice Guidelines. 4/1/11.*
5. *Chou R. Subacute and chronic low back pain: Nonsurgical interventional treatment. In: UpToDate Online Journal [serial online]. Waltham, MA: UpToDate; updated June 10, 2021. Literature review current through October 2023.*
6. *Manchikanti L, Singh V, Falco FJ, et al. Lumbar facet joint nerve blocks in managing chronic facet joint pain: one-year follow-up of a randomized, double-blind controlled trial: Clinical Trial NCT00355914. Pain Physician. 2008 Mar-Apr; 11(2):121-32.*
7. *Novitas Solutions Local Coverage Determination (LCD) L34892: Facet Joint Interventions for Pain Management.*
8. *Sehgal N, Dunbar BE, Shah RV, Colson J. Systematic review of diagnostic utility of facet (zygapophyseal) joint injections in chronic spinal pain: an update. Pain Physician 2007; 10(1): 213-28.*
9. *Adbi, Salahadin. Complex regional pain syndrome in adults: Treatment, prognosis, and prevention. In: UpToDate Online Journal [serial online]. Waltham, MA: UpToDate; updated July 1, 2022.*
10. *Speed CA. Injection therapies for soft-tissue lesions. Best Pract Res Clin Rheumatol 2007; 21 (2): 333-47.*
11. *Staal JB, de Bie R, de Vet HC, Hildebrandt J, Nelemans P. Injection therapy for subacute and chronic low-back pain. Cochrane Database Syst Rev. 2008 Jul 16 ;(3):CD001824.*
12. *North American Spine Society (NASS). Evidence-based clinical guidelines for multidisciplinary spine care: Diagnosis and treatment of low back pain. 2020*

MEDICAL POLICY

POLICY TITLE	EPIDURAL STEROID INJECTIONS FOR BACK PAIN AND FACET NERVE BLOCKS
POLICY NUMBER	MP 4.014

13. Won HS, Yang M, Kim YD. Facet joint injections for management of low back pain: a clinically focused review. *Anesth Pain Med (Seoul)*. 2020;15(1):8-18. doi:10.17085/apm.2020.15.1.8
14. Cohen SP, Moon JY, Brummett CM, White RL, Larkin TM. Medial Branch Blocks or Intra-Articular Injections as a Prognostic Tool Before Lumbar Facet Radiofrequency Denervation: A Multicenter, Case-Control Study. *Reg Anesth Pain Med*. 2015;40(4):376-383. doi:10.1097/AAP.0000000000000229
15. Cohen SP, Bhaskar A, Bhatia A, et al. Consensus practice guidelines on interventions for lumbar facet joint pain from a multispecialty, international working group. *Reg Anesth Pain Med*. 2020;45(6):424-467. doi:10.1136/rapm-2019-101243

X. POLICY HISTORY

[Top](#)

MP 4.014	02/12/2020 Consensus review. Policy statement unchanged. References updated
	11/17/2020 Administrative update. Codes 0228T, 0229T, 0230T and 0231T deleted. Effective 1/1/2021.
	01/08/2021 Consensus review. Policy statement unchanged. References updated. Coding reviewed.
	09/07/2021: Administrative update. Addition of new ICD-10 codes. Effective date 10/1/2021.
	11/22/2022 Consensus review. For clarity, changed statement section heading from “Facet Joint Nerve Block Injections” to “Facet Joint Injections/Medial Branch Blocks”. No change to intent or coverage. Updated cross-references, FEP, Background, Rationale, and references. Updated coding table.
	12/01/2023 Minor review. To facet joint/medial branch block, separated out sections with updated criteria for initial diagnostic injection, second diagnostic injection, and therapeutic injections. To epidural section, clarified which approach is eligible for multi-levels and updated time between injections as 14 days instead of 30 days. Updated policy guidelines and references. No changes to coding.

[Top](#)

Health care benefit programs issued or administered by Capital Blue Cross and/or its subsidiaries, Capital Advantage Insurance Company[®], Capital Advantage Assurance Company[®] and Keystone Health Plan[®] Central. Independent licensees of the BlueCross BlueShield Association. Communications issued by Capital Blue Cross in its capacity as administrator of programs and provider relations for all companies.